



PATIENT NAME

PATIENT ID

PATIENT DOB

## Remdesivir and TACTIC-R Trial

Treatment of patients with Remdesivir is not a contraindication to participation in TACTIC-R, indeed its use is encouraged as standard of care, if the patient is eligible.

To check if your patient is eligible for Remdesivir, TACTIC-R, or both:

1. **Must be a hospital in-patient, not receiving invasive mechanical ventilation, ECMO, or cardiovascular support (vasopressors or inotropes)**
2. **Complete the ELIGIBILITY CRITERIA TICK BOX CHARTS**

If all boxes are ticked for Remdesivir, then refer to respiratory to ensure there are no contraindications to treatment.

If all boxes are ticked for TACTIC-R, then refer to the TACTIC-R team to ensure there are no contraindications to treatment.

If all boxes are ticked for both Remdesivir and TACTIC-R, then your patient can be considered for **both**.

\_\_\_\_\_ Sign and Date



PATIENT NAME
PATIENT ID
PATIENT DOB

**Prescreening Checklists for Determining if a Patient is Eligible for Remdesivir and/or TACTIC-R**

<b>Eligibility Criteria for Remdesivir</b>	<b>TICK if YES</b>
Age over 12 years	
Suspected or laboratory confirmed SARS-CoV-2 infection	
Weight ≥40kg	
Oxygen saturations ≤94% on room air OR Requiring supplemental oxygen OR Requiring non-invasive or invasive ventilation or ECMO	
<b>ONLY if ALL boxes ticked, discuss with Infectious Diseases or Respiratory consultant (according to Trust policy) as may be suitable for Remdesivir</b>	

<b>Eligibility Criteria for TACTIC-R Trial</b>	<b>TICK if YES</b>
Age over 18 years	
Clinical picture strongly suggestive of COVID, <b>with or without</b> a positive Covid-19 test	
TACTIC risk score of 4 or more (if radiographic severity score 3 or less), or 3 or more (if radiographic severity score 4 or more). See chart overleaf.	
Not pregnant or breastfeeding	
Not participating in RECOVERY or REMAP-CAP	
Not currently being treated with probenecid, chronic IVIG, Baricitinib or Ravulizumab	
Not known to have active TB, unresolved Neisseria meningitidis infection, active Hepatitis B/C or Varicella Zoster (no blood testing required to exclude these)	
Creatinine clearance ≥30 ml/min	
ALT ≤250 U/L	
VTE prophylaxis or maintenance therapy not contraindicated	
<b>ONLY if all boxes ticked, may be suitable for TACTIC-R, please discuss with TACTIC-R team</b>	



PATIENT NAME

PATIENT ID

PATIENT DOB

### TACTIC RISK SCORE

(needed for for TACTIC-R)

	Score (1 point each)
Radiographic severity score $\geq 3$ (see below)	
Male gender	
Non-white ethnicity	
Diabetes	
Hypertension	
Neutrophils $> 8.0$	
Age $> 40$	
CRP $> 40$	
<b>TOTAL</b>	

### RADIOGRAPHIC SEVERITY SCORE

(check CXR or CT chest scout images or D/W respiratory)

Score 0-8.  
Score each lung separately.

0 = normal  
1 =  $\leq 25\%$  infiltrate  
2 = 25-50% infiltrate  
3 = 50-75% infiltrate  
4 =  $\geq 75\%$  infiltrate

**All TACTIC-R patients must have a TACTIC risk score of 4 or more (if radiographic severity score 3 or less), or 3 or more (if radiographic severity score 4 or more).**

(Online training for performing Radiographic Severity Score:  
[https://www.camcovidtrials.net/trials/view,tacticr\\_1.htm](https://www.camcovidtrials.net/trials/view,tacticr_1.htm))